



UFRJ



FROM PARTICLES TO FIELDS:

HOW THE cDFT AND PF MODELS CAN IMPROVE THE FLUID-SOLID DESCRIPTION?

Elvis do A. Soares, PhD.

elvis.asoares@gmail.com




EQUIFASE
Brasil 2022

XII Equifase, Campinas-SP, October 24-27, 2022

Do You Have 10 MINUTES TO
TALK ABOUT CLASSICAL DFT?

A BRIDGE BETWEEN DIFFERENT LENGTH SCALES

MC, MD



Individual particles

**Approximation by a
Free-energy functional**




DFT



Density field

MULTI-SCALE MODELLING



CLASSICAL DENSITY FUNCTIONAL THEORY

Grand Potential

$$\Omega(V, T, \mu) = F[\rho(\mathbf{r})] + \int_V d\mathbf{r} [V_{\text{ext}}(\mathbf{r}) - \mu] \rho(\mathbf{r})$$

Equilibrium

$$\Omega[\rho(\mathbf{r})] \geq \Omega[\rho(\mathbf{r})]_{\text{eq.}}$$

Equilibrium Condition

$$\left. \frac{\delta \Omega}{\delta \rho(\mathbf{r})} \right|_{V, \mu, T} = k_B T \ln(\Lambda^3 \rho(\mathbf{r})) + \frac{\delta F_{\text{exc}}[\rho]}{\delta \rho(\mathbf{r})} + V_{\text{ext}}(\mathbf{r}) - \mu = 0$$

Free Energy Functional

$$F[\rho] = F_{\text{id}}[\rho] + F_{\text{exc}}[\rho]$$





Excess Functionals

FMT, MFA,
WDA, MMFA,
FMSA, ...

Particle-particle
interaction

External potential acts
as a chemical potential


LENNARD-JONES FLUIDS





Lennard-Jones potential

$$u(r) = 4\epsilon \left[\left(\frac{\sigma}{r} \right)^{12} - \left(\frac{\sigma}{r} \right)^6 \right]$$


RADIAL DISTRIBUTION FUNCTION



Data from X-Ray
Experiment





RADIAL DISTRIBUTION FUNCTION




liquid

$$\rho(\mathbf{r}) = \rho_b e^{-\beta V_{\text{ext}}(\mathbf{r})}$$

Data from X-Ray
Experiment





RADIAL DISTRIBUTION FUNCTION



liquid

Data from X-Ray
Experiment







$$\rho(\mathbf{r}) = \rho_b e^{-\beta V_{\text{ext}}(\mathbf{r}) - \beta \frac{\delta F_{\text{exc}}[\rho]}{\delta \rho(\mathbf{r})} + \beta \mu_{\text{exc}}}$$



Density
Correlations are
very important!

Non-local DFT


FLUID INSIDE A SLITLIKE PORE



Snook, I. K., & van Megen, W. (1980). The Journal of Chemical Physics, 72(5), 2907-2913




Tang, Y., & Wu, J. (2004). Physical Review E, 70(1), 011201.

ADSORPTION IN ACTIVATED CARBON



PCP-SAFT


$$F_{\text{exc}} = F_{\text{hc}} + F_{\text{disp}} + F_{\text{qq}}$$



CAN WE GO BEYOND SYMMETRICAL
GEOMETRIES?

DFT IN 3D GEOMETRIES

$$\Omega[\{\rho_{\mathbf{R}}\}] = \sum_{\mathbf{R}} \left[k_B T \rho_{\mathbf{R}} (\ln(\rho_{\mathbf{R}} \sigma^3) - 1) + F_{\mathbf{R}}^{\text{exc}} + \rho_{\mathbf{R}} V_{\mathbf{R}}^{\text{exc}} - \mu \rho_{\mathbf{R}} \right] \Delta^3$$




Convolution (FFT)

$$n(\mathbf{r}) = \int \rho(\mathbf{r}') \omega(\mathbf{r} - \mathbf{r}') d\mathbf{r} \implies \tilde{n}_{\mathbf{k}} = \tilde{\rho}_{\mathbf{k}} \tilde{\omega}_{\mathbf{k}}$$

Extremum condition (FIRE)


$$\left\| \frac{\partial \Omega}{\partial \rho_{\mathbf{R}}} \right\| < \epsilon_{\text{tol}}$$

GRIDSIZE VS. TIME CONSUMPTION




Grid	Time(s)
0.01 ¹	8

GRIDSIZE VS. TIME CONSUMPTION




Grid	Time(s)
0.01 ¹	8
0.05 ³	545

GRIDSIZE VS. TIME CONSUMPTION



Grid	Time(s)
0.01^1	8
0.05^3	545
0.1^3	177


GRIDSIZE VS. TIME CONSUMPTION



Grid	Time(s)
0.01^1	8
0.05^3	545
0.1^3	177
0.2^3	27

METAL ORGANIC FRAMEWORK

MOF-5



DREIDING Force Field
H, C, O, Zn

External Potential



$$V^{\text{ext}}(\mathbf{r}) = \sum_{j \in \text{solid}} u_j^{(\text{lj})}(|\mathbf{r} - \mathbf{r}_j|)$$

Lorentz-Berthelot mixing rules


$$\sigma_{ij} = \frac{1}{2}(\sigma_{ii} + \sigma_{jj})$$

$$\epsilon_{ij} = (\epsilon_{ii}\epsilon_{jj})^{1/2}$$

EQUATION OF STATE – MBWR – LJ FLUID





EXTERNAL POTENTIAL FOR H_2



ADSORPTION ISOTHERMS

No fitted
parameter!


○ Exp.Data
— DFT




ISOSURFACES OF H_2 LOCAL DENSITY

$P = 60 \text{ bar}$
 $T = 300 \text{ K}$

$\rho = 0.001 \text{ \AA}^{-3}$




$\rho = 0.02 \text{ \AA}^{-3}$




ISOSURFACES OF CH_4 LOCAL DENSITY

$P = 60 \text{ bar}$
 $T = 300 \text{ K}$

$\rho = 0.001 \text{ \AA}^{-3}$




$\rho = 0.02 \text{ \AA}^{-3}$



VIDEO

$$P = 60 \text{ bar}$$
$$T = 300 \text{ K}$$


$$\rho = 0.08 \text{ \AA}^{-3}$$



CONCLUSIONS AND PERSPECTIVES

CONCLUSIONS

- The **size correlations are very important** to describe the high concentration systems;
- The **fluid-solid interaction can be described** by the DFT on the same level of MC/MD;
- There is a connection between **DFT and Phase-Field Models. (Possible Scale Integration)**
- **Future works:**
 - Extensions to associative interactions, chain-like systems.
 - Quantum DFT? Chemisorption
 - Machine-learning functionals



ACKNOWLEDGMENTS



elvis.asoares@gmail.com



Elvis Do Amaral Soares



elvissoares



PETROBRAS



instagram.com/atoms.ufrj



facebook.com/atoms.ufrj



youtube.com/c/atoms-ufrj



FEDERAL UNIVERSITY OF RIO DE JANEIRO - UFRJ

**Applied Thermodynamics and Molecular Simulation
ATOMS**

Av. Athos da Silveira, nº 149, Centro de Tecnologia
Cidade Universitária, Rio de Janeiro, RJ, Brasil

Postal Code: 21941-909 / Phone Number: +55 21 3938 7428